


Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.



Chapter 6 Cranberry

Catherine C. Neto and Joe A. Vinson.

6.1. CRANBERRY: INTRODUCTION AND TRADITIONAL ORIGINS

The name “cranberry” reportedly derives from the Pilgrim name for the fruit “craneberry,” because the small, pink blossoms that appear in the spring resemble the head and bill of a Sandhill crane. Cranberries are unique among fruits. They can grow and survive only under a special combination of factors. They require acid peat soil, adequate freshwater supply, sand, and a growing season that stretches from April to November and is followed by a period of dormancy in the winter months that provides an extended cold period, necessary for fruiting buds to mature. Contrary to popular belief, cranberries do not grow in water. Instead, they grow on vines in impermeable beds that are layered with sand, peat, gravel, and clay. These beds, commonly known as “bogs,” were originally made by glacial deposits, but now, they can be made by humans. Growers do not have to replant, since an undamaged cranberry vine will survive indefinitely. Some vines in Massachusetts are more than 150 years old. European settlers adopted the Native American uses of the fruit and found that cranberry is a valuable bartering tool. American whalers and mariners carried cranberries on their voyages to prevent scurvy. In 1816, Captain Henry Hall was the first to successfully cultivate cranberries ([Cape Cod Cranberry Grower’s Association](http://www.cranberries.org/cranberries/history.html), <http://www.cranberries.org/cranberries/history.html>). Today, cranberries are available in both fresh and processed forms. There are many varieties of cranberry fruit; white cranberry is an early harvest cranberry, which is picked 2–3 weeks prior to ripening.

There are two major species of cranberry: the American cranberry (*Vaccinium macrocarpon*) and the European cranberry (*V. oxycoccos*). The European cranberry fruit is smaller (0.6–1.2 cm) and only half the size of the American fruit. The American cranberry, which is frequently cultivated, is a member of the Ericaceae family, evergreens, creeping shrubs native to the cool, temperate, acidic soils and peat wetlands of Northeastern United States and southern Canada. Latvia, Belarus, Azerbaijan, and Ukraine are other cranberry-producing countries in Europe, with Turkey just beginning cranberry cultivation. The United States and Canada together account for more than 90% of the world’s production ([Zhao 2007](#)). The forecast for U.S. cranberry production in 2010 is 735 million pounds up by 6% from 2009 figures and the second highest

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

yearly production. Wisconsin is expected to lead all U.S. states in the production of cranberries, with 435 million pounds, followed by Massachusetts, with 195 million. New Jersey, Oregon, and Washington are also expected to have substantial production (Cranberries, National Agricultural Statistics Service (NASS), Agriculture Statistics Board, United States Department of Agriculture (USDA), August 17, 2010).

Cranberries are low-growing, woody, perennial vines with small, alternate, and ovate leaves. The plant produces stolons (horizontal stems) having a height of up to 6 feet (2 m). Short, vertical branches, or uprights, 2–8 inches (5–20 cm) in height, grow from buds on the stolons, and these can be either vegetative or fruiting. Each fruiting upright may contain as much as seven flowers. Pollination is primarily via domestic honeybees (Cranberry Institute, East Wareham, Massachusetts). Cranberries were first used by Native Americans, who discovered the wild berry's versatility. Native Americans used cranberries in a variety of foods, the most popular being pemmican, a high-protein combination of crushed cranberries, dried deer meat, and melted fat. They also used it as a medicine to treat arrow wounds and as a dye for rugs and blankets.

[Go to:](#)

6.2. PRODUCTION AND CONSUMPTION

About 95% of the cranberries cultivated are processed into products such as juice drinks, sauce, and sweetened, dried cranberries. The remaining 5% are sold fresh to consumers. Cranberries used for processing are commonly frozen in bulk containers shortly after arriving at a receiving station. The primary method of harvesting cranberries takes advantage of the ability of cranberries to float. In this method, which is called wet harvesting, cranberry fields are flooded with water. After flooding, eggbeater-like devices stir up the water with sufficient force to dislodge the berries from the vines. When the berries float on the surface of water, they are pulled to the shore with hinged two-by-fours and loaded in trucks for delivery to the factory. Wet-harvested cranberries are generally processed for juice and sauce, because once a berry gets wet, there is increased chance of spoilage unless it is processed rapidly. Dry harvesting is the second method of harvesting cranberries. Because fully ripe berries are easily dislodged from their vines, this method employs a lawn mower-type machine that combs the berries from the vines. On arrival at the factory, dry-harvested cranberries are subjected to a unique test. The superior berries are sorted from those that are bruised, soft, or rotten by taking advantage of the fact that berries bounce when they are of good quality. Along a conveyor belt, each berry must successfully bounce over a series of wooden barriers. Berries that fail this simple test fall into a disposal bin. The ones that pass the bounce test are sold as fresh fruit, typically packaged in plastic (MotherLindas.com, <http://www.motherlindas.com/cranberries.htm>).

There are approximately 450 cranberries in 1 pound, 4,400 cranberries in 1 gallon of juice, and 45,000 cranberries in a 100-pound barrel. It takes about 200 cranberries to make one can of cranberry sauce ([foodreference.com](http://www.foodreference.com/html/fcranberries.html), <http://www.foodreference.com/html/fcranberries.html>). The cranberry is an underconsumed fruit in the United States. Consumption of different types of fruits in the United States is shown in [Figure 6.1](#). Americans eat on an average 326 g of fruit per day, and only 5.3% of this intake is berries (18 g). This translates to 2.8 g or about three cranberries per day. This is equivalent to drinking 0.25 mL of cranberry juice per day. The dominant cranberry product consumed in the United States is juice. In comparison, the per capita fruit consumption in the European Union in 2007 was considerably less, only 266 g/day

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

([Freshfel Europe 2009](#)). Average per capita consumption of berries in Norway was only 5 g/day ([Ellingsen et al. 2008](#)). Thus, Europeans on the average consume fewer berries than the U.S. population.




FIGURE 6.1

Per capita consumption of fruits in the United States in 2007. (From USDA/Economic Research Service. Food Availability (Per Capita) Data System. <http://www.ers.usda.gov/Data/FoodConsumption>. Data last updated February 16, 2010. With permission.)

[Go to:](#)

6.3. PHYTOCHEMICAL COMPOSITION

It is interesting to discover via a PubMed search that there are over 52,000 references for flavonoids (<http://www.ncbi.nlm.nih.gov/sites/entrez>, accessed Nov. 14, 2010). Among the subclasses, the order of references is flavonols > flavonones > anthocyanins > flavanols (catechins). [Table 6.1](#) lists the potentially bioactive compounds identified and often quantified in cranberries. [Table 6.2](#) lists the flavonoids quantified in cranberry as listed in the U.S. Department of Agriculture (USDA) database. Representative structures of the major constituents are shown in [Figures 6.2](#) and [6.3](#).

Class of Compounds
Anthocyanins
Cyanidin

TABLE 6.1

Potentially Bioactive compounds Present in Cranberries and Cranberry Juice.

Flavonoid	Milligrams per 100 Gra
Flavonols, total ^c	
Quercetin ^a	15.6
Myricetin ^a	6.7
Kaempferol ^a	0.0
Anthocyanins, total ^c	
Cyanidin ^a	41.4
Peonidin ^a	42.1

TABLE 6.2

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

Content of Flavonoids in Cranberry Fruit by Flavonoid Class, as Reported by the USDA Databases on Nutrient Composition of Foods.



FIGURE 6.2

Structures of some of the major anthocyanins (cyanidin and peonidin), flavonols (quercetin and myricetin), phenolic and organic acids (*p*-coumaric acid, chlorogenic acid, and benzoic acid), flavan-3-ols (epicatechin), and terpenoids (ursolic acid) in cranberry ([more...](#))




FIGURE 6.3

Cranberry uniquely contains A-type linkage (left), whereas other foods contain B-type linkage (right).

6.3.1. ANTHOCYANINS

When one thinks of cranberries, it is the color red that comes to mind. It is due to the presence of anthocyanins, which have been extensively studied for health benefits. There are six aglycones in the anthocyanin class, and the number of potential isomers is extremely large due to different positions of sugar attachment and the large number of different mono- and disaccharides (glycosides) and, the less common, but possible, phenolic acids and acyl groups that can be attached. Anthocyanin content has been reported to be as high as 91.5 mg/100 g ripe fruit at harvest depending on the cultivar ([Wang and Stretch 2001](#); [Wu et al. 2006](#)). Fruit of the early black cultivar tends to contain more anthocyanins and proanthocyanidins (PACs) than other cranberry cultivars ([Wang and Stretch 2001](#); [Vorsa and Howell 2003](#)). Cranberry, compared to many other berries, has a very small number of anthocyanin isomers ($n = 13$), the major ones being galactosides and arabinosides of cyanidin and peonidin. A very similar anthocyanin profile exists in cranberry juice ([Fuleki and Francis 1968](#); [Ohnishi et al. 2006](#)). Only the most recent references are listed in [Table 6.1](#), since the newer technology liquid chromatography-mass spectrometry (LC-MS)/MS is better able to unequivocally identify compounds in mixtures than the classical methods of column chromatography followed by crystallization and spectral investigation. Whole cranberries contain a very broad concentration range of anthocyanins, 180–596 mg/kg fresh weight (fw; [Bilyk and Sapers 1986](#)), which also varies according to maturity from 0.8 to 111.0 mg/kg fw from the green to dark red stage ([Celik et al. 2008](#)). Cranberry juice

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

contains much lower levels of anthocyanins, ranging from 1.3 mg/100 mL ([Prior et al. 2001](#)) to 2.5 mg/100 mL of juice ([Cunningham et al. 2003](#)).

6.3.2. FLAVONOLS

The total amount of flavonols in cranberry ranges from 200 to 400 mg/kg ([Zheng and Wang 2003](#); [Cunningham et al. 2003](#); [Vvedenskaya et al. 2004](#); [Neto 2007b](#)). Cranberry is the best source of flavonols among 30 flavonol-containing plant foods studied ([Aherne and O'Brien 2002](#)), and the flavonol content of cranberry is almost twice as high as 12 other commonly consumed fruit juices, including pomegranate and grape ([Mullen, Marks, and Crozier 2007](#)). Quercetin is the most abundant flavonol in cranberry, and it varies from 11 to 25 mg/100 g, primarily as the 3-*o*-galactoside ([Yan et al. 2002](#); [Vvedenskaya et al. 2004](#)). Cranberry is also the best source of quercetin ([Manach et al. 2004](#)). Myricetin is the second most abundant flavonol, followed by kaempferol ([Vvedenskaya et al. 2004](#)). These compounds are yellow in color, and there are 20 different flavonol glycosides in cranberry, as confirmed by another article ([Vvedenskaya and Vorsa 2004](#)).

6.3.3. FLAVAN-3-OLS (CATECHINS)

Total catechins in raw cranberry average 17 mg/100 g, with epicatechin being the most abundant ([Harnly et al. 2006](#)). Berries have the highest catechin content among fruits. Cranberries have the highest levels of monomers (catechin + epicatechin) among berries, and monomer levels in cranberries are twice as high as those in blueberries ([Gu et al. 2004](#)). Cranberry juice contains 6 mg/l of catechins ([Gu et al. 2004](#)). One liter of cranberry juice has catechin and epicatechin monomers as 100 g of dark chocolate.

6.3.4. PROANTHOCYANIDINS

These compounds are the largest class of potential bioactives in cranberries. The PACs are also commonly called condensed tannins or poly flavan-3-ols, and are responsible for the bitter astringent taste of cranberries due to the binding of saliva proteins. Cranberries have the most dimers, trimers, 4–6mers, and 7–10mers compared with any other fruit studied ([Gu et al. 2004](#)). The most unique aspect of cranberry polyphenols is the occurrence of A-type linkages (C2→O→C7) between epicatechin units, as shown in [Figure 6.3](#) ([Foo et al. 2000b](#)). These linkages are found elsewhere only in peanuts, blueberries, and plums ([Gu et al. 2003](#)). The average concentration of PACs is 419 mg/100 g (0.4%) by weight in cranberries and 231 mg/l in cranberry juice ([Gu et al. 2004](#)). These compounds exhibit potent in vitro antiadhesion bioactivity and are present in much higher concentrations in cranberries than in other foods.

6.3.5. BENZOIC AND PHENOLIC ACIDS

Benzoic and phenolic acids represent 0.57% of the weight of fresh cranberries ([Zuo, Wang, and Zhan 2002](#)). Benzoic acid forms 80% of the total organic acids contained in cranberry juice. There exist 14 other benzoic and phenolic acids in cranberry juice, and *p*-coumaric acid is the most prevalent hydroxycinnamic acid. Many of these acids are bound to glucose and polysaccharides in cranberry ([He and Liu 2006](#)).

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

6.3.6. NONFLAVONOID POLYPHENOLS

Phloridzin is the most prevalent member of the class of dihydrochalcones present in cranberries at 120 mg/kg ([Turner et al. 2007](#)). This compound is present in very high quantities in apple, and is a dihydrochalcone with glucose-lowering activity in an animal model of diabetes ([Masumoto et al. 2009](#)). Resveratrols, one of the most studied polyphenolic compounds, with 2937 PubMed references, are found in cranberry juice at extremely low levels of 0.2 mg/l ([Wang et al. 2002](#)). This is much lower than the resveratrol content of red wine (14 mg/l; [Baur and Sinclair 2006](#)). Other compounds belonging to this stilbene class are present at very low levels in cranberry juice, and thus are probably nonbioactive.

6.3.7. TERPENES AND STEROLS

Terpenes include the volatile compounds that are responsible for the flavor and aroma of cranberries ([Croteau and Fagerson 1968](#)), as well as larger oily or waxy compounds. Cranberry fruit contains the triterpenoid ursolic acid (UA) in its peel ([Figure 6.3](#)), existing in aglycone form as well as in *cis* and *trans* *p*-hydroxycinnamate esters ([Murphy et al. 2003](#)). Quantitative analysis of cranberry fruit and products by LC-MS has determined the UA content of the whole cranberry fruit of different cultivars to be between 60 and 110 mg per 100 g of fresh fruit ([Kondo 2006](#)). UA has been reported to be a constituent of other fruits, including apples and highbush blueberries ([Wang et al. 2000](#)). The iridoid glycosides listed in [Table 6.1](#) are believed to be unique to the *Vaccinium* species. Cranberries also contain the carotenoid lutein, as well as other carotenoids in lesser quantities. It is worth noting that although the compounds listed in [Table 6.2](#) are the result of studies conducted on the American cranberry (*V. macrocarpon*), a similar variety of anthocyanins, flavonols, nonflavonoid polyphenols, and phenolic acids are found in the European cranberry (*V. oxycoccus*) even though this variety has not been studied as extensively as its American counterpart ([Häkkinen et al. 1999](#); [Kähkönen, Hopia, and Heinonen 2001](#)).

[Go to:](#)

6.4. BIOAVAILABILITY OF CRANBERRY PHYTOCHEMICALS

The first evidence of human absorption of cranberry compounds was demonstrated by a single subject who consumed 1800 mL of 27% commercial cranberry juice after an overnight fast ([Zhang and Zuo 2004](#)). The plasma was then extracted and derivatized prior to assay. Sixteen compounds were identified in plasma by GC-MS from blood samples drawn 45 and 270 minutes after consumption. After 45 minutes, five acids were identified: (1) benzoic acid, (2) *o*-hydroxybenzoic acid (salicylic acid), (3) *p*-hydroxyphenylacetic acid, (4) 2,3-dihydroxybenzoic acid, and (5) 2,4-dihydroxybenzoic acid. Phenylacetic acid and dihydroxybenzoic acid are not found in cranberry juice and, thus, they are the products of polyphenol breakdown or bacterial metabolism in the gut. At 270 minutes, the same five acids and ferulic acid plus sinapic acid were found. Anthocyanins were not analyzed in the study. In 2005, an early-morning urinary excretion study was conducted with 11 females who were administered a commercial cranberry juice three times a day (250 mL each) for 2 weeks ([Duthie et al. 2005](#)). High-performance liquid chromatography (HPLC) with electrochemical detection was used after urine hydrolysis with NaOH and extraction. After 1–2 weeks, the urinary level of salicylic acid (metabolite of

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

salicylic acid) significantly increased. This compound was also increased in the fasting plasma after 2 weeks of consumption of cranberry juice. A Japanese group gave 200 mL of 100% cranberry juice to 11 subjects, and urine was collected before administering the juice and again 24 hours afterwards ([Ohnishi et al. 2006](#)). Six of the 12 anthocyanin glycosides identified in cranberry juice by LC-MS/MS were found in the collected urine. Peonidin 3-*o*-galactoside, the second most plentiful anthocyanin in the juice, was the major anthocyanin in the urine. Over 5% of the consumed anthocyanins were found in the urine. This level of absorption is at least 10 times higher than that found for other berry juices, which ranges from 0.08% for blood orange juice ([Giordano et al. 2007](#)) to 0.2% for red wine and red grape juice ([Frank et al. 2003](#)) and 0.4% for black currant juice ([Bitsch et al. 2004](#)). A safety study in which a cranberry powder was given to 65 healthy women for 8 weeks at a dose of 1200 mg/day and the effects were compared with a placebo ([Valentova et al. 2007](#)) showed no toxicity for the powder. In decreasing order of content, hippuric acid, salicylic acid, quercetin glucuronide, and dihydroxybenzoic acid isomers were found in the urine by LC-MS.

[Go to:](#)

6.5. HUMAN STUDIES RELEVANT TO HEART DISEASE AND DIABETES

Growing evidence points to many potential pharmacological activities for cranberry and cranberry juice. Due to its high content of flavonoids and phenolic acids, cranberry ranks highly among fruits in both antioxidant quality and quantity ([Vinson et al. 2001](#)). In fact, cranberry has the highest level of fw polyphenols among a group of 22 fruits studied. These antioxidant properties are likely to contribute to cranberry's disease-fighting ability.

Cranberry compounds can cause an improvement in antioxidant status, which might be beneficial with respect to chronic diseases. The potential cardiovascular benefits have been reviewed by two groups of researchers ([McKay and Blumberg 2007](#); [Ruel and Couillard 2007](#)). A single dose of 500 mL 27% cranberry administered to nine healthy women increased plasma antioxidant capacity against a sucrose control ([Pedersen et al. 2000](#)). The authors stated that this increase was due to the presence of ascorbic acid (vitamin C) in cranberry juice. Unfortunately, ascorbic acid was not included in the control sucrose solution. Another group found that cranberry juice relative to a control containing sugars and ascorbic acid significantly raised the plasma antioxidant capacity in normal subjects ([Vinson et al. 2008](#)). They also found that when cranberry juice was spiked on human plasma, the lipoproteins low-density lipoprotein (LDL) and very-low-density lipoprotein (VLDL) were protected from oxidation in a dose-dependent manner. This second effect is directly related to atherosclerosis ([Steinberg 2009](#)).

The proof of benefit for disease prevention should be shown by a human supplementation study. There have been a few of these studies for cranberry, all done in the last 6 years. Cranberry powder equivalent to 240 mL of cranberry juice per day was given to 12 diabetic men and 15 diabetic women for 12 weeks ([Chambers and Camire 2003](#)). There was no effect on glucose control. According to the authors, no change was observed on glucose control because the basic cranberry powder might have undergone processing in its production from cranberry juice. In 2008, a Japanese group reported the effect of a cranberry powder and a placebo given to 30 type 2 diabetic subjects for 12 weeks ([Lee et al. 2008](#)). There was no effect of cranberry on fasting glucose or glycated hemoglobin levels. However, it was found that total cholesterol and LDL significantly decreased and the cholesterol/high-density lipoprotein (HDL) ratio significantly

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

increased in the cranberry group. All the subjects were taking oral hypoglycemic drugs and, thus, their plasma glucose levels were under control. In another study, type 2 diabetics ($n = 12$) were given a low-calorie cranberry juice at a dose of 240 mL ([Wilson et al. 2008](#)). The juice was found to give a more favorable glucose and insulin response than a sugared water control.

Cranberry juice was given to 20 women at a dose of 750 mL/day for 2 weeks ([Duthie et al. 2006](#)). No effect on plasma lipids, antioxidant enzyme, or DNA oxidation was found, but plasma ascorbic acid was increased. Healthy men ($n = 21$) were given 7 mL/kg of cranberry juice for 14 days ([Ruel et al. 2005](#)). Although there was no effect on lipids, oxidized LDL was significantly decreased (10%) and plasma antioxidant capacity was significantly increased (7%) in the subjects. Inflammatory markers were not improved. In the latest study by the same group, men with elevated LDL and body mass index (BMI) were given low-calorie cranberry juice at doses of 125, 250, and 500 mL/day for three successive 4-week periods ([Ruel et al. 2007](#)). Significant decreases in oxidized LDL, systolic pressure, and the inflammatory markers ICAM-1 and VCAM-1 were recorded at the end of the study. All cranberry effects in this study were found to be beneficial with respect to heart disease.

[Go to:](#)

6.6. INVESTIGATION OF CRANBERRY CONSTITUENTS THAT CONTRIBUTE TO IN VITRO ANTICANCER ACTIVITY

Cranberry's role as a potential chemopreventive agent is gradually emerging from in vitro model studies by various researchers. Cranberry compounds may act against cancers by inhibiting oxidative stress or by other pathways. Over the past 10 years, numerous in vitro studies have appeared addressing the effects of cranberry and its constituents against tumor cell proliferation and the possible mechanisms of action. Several reports of in vivo studies have appeared recently, lending support to the tumor-inhibiting potential of cranberry. These studies are discussed. A summary of in vitro and in vivo studies on the anticancer properties of cranberry, the constituents contributing to these properties, and some possible mechanisms of action are discussed.

The first in vitro report of anticancer activity in *Vaccinium* fruit appeared in 1996. A University of Illinois study reported that extracts of cranberry, bilberry, and other species inhibit ornithine decarboxylase (ODC) expression and induce the xenobiotic detoxification enzyme quinone reductase in vitro ([Bomser et al. 1996](#)). Some Canadian researchers reported that cranberry juice extracts inhibit breast tumor cell growth ([Guthrie 2003](#)), and later showed that an extract of cranberry press cake inhibits proliferation of MCF-7 and MDA-MB-435 breast cancer cells ([Ferguson et al. 2004](#)). Later studies by various researchers have focused on identifying the anticancer constituents of cranberry, which fall within several possible classes, including triterpenoids and flavonoids.

6.6.1. URSOLIC ACID AND DERIVATIVES

A bioassay-guided fractionation approach was used to examine in vitro the antitumor activities of whole cranberry fruit and juice, extracts and fractions, and finally individual compounds or subfractions within structural classes. An ethyl-acetate extract of whole cranberry fruit inhibited growth of human tumor cell lines in vitro ([Yan et al. 2002](#)). From ethyl-acetate-soluble extracts, two hydroxycinnamate esters of UA were isolated and identified that inhibited the growth of

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

several types of human tumor cells in vitro, including MCF-7 breast, HT-29 colon, DU145 prostate, H460 lung, ME180 cervical epidermoid, and K562 leukemia cell lines ([Murphy et al. 2003](#)). Growth-inhibitory concentrations (GI_{50}) for esters were 11–28 $\mu\text{g/mL}$, depending on the cell line. LC-MS analysis of cranberry fruit found that in addition to UA, its hydroxycinnamate esters are present in whole cranberry fruit in quantities averaging about 15–20 mg per 100 g of fresh fruit ([Kondo 2006](#)). UA isolated from cranberry fruit was also reported to inhibit the proliferation of HepG2 human liver cancer cells and MCF-7 cells ([He and Liu 2006](#)). It was also found to inhibit tumor colony formation in a dose-dependent manner in HT-29 and HCT116 colon tumor models over a 2-week period, based on clonogenic assays ([Liberty, Hart, and Neto 2007](#)).

UA has been reported as a constituent of herbal medicines marketed worldwide for treating inflammatory conditions ([Kim et al. 2004](#)), but bioavailability data on this compound is lacking. In vivo cancer studies of UA are scarce, but one mouse model study reported that a UA dose of 100 mg/kg inhibits murine fibrosarcoma FSaII growth ([Lee et al. 2001](#)). Numerous reports have appeared on the in vitro antitumor activity of UA ([Novotny, Vachalkova, and Biggs 2001](#)), and these suggest a variety of mechanisms of action, including early G1 cytostatic effect ([Es-Saady et al. 1996](#)), induction of apoptosis ([Baek et al. 1997](#)), enhancement of intracellular Ca^{2+} signaling ([Novotny, Vachalkova, and Biggs 2001](#)), enhanced release of cytochrome *c*, caspase activation ([Andersson et al. 2003](#)) and downregulation of inhibitor of apoptosis proteins (c-IAPs; [Choi et al. 2000](#)), increased expression of $p21_{\text{WAF1}}$ ([Kim et al. 2000](#)), and decreased matrix metalloproteinase (MMP)-9 expression ([Cha et al. 1996](#)). In addition to inhibiting the growth of several tumor cell lines, the hydroxycinnamoyl esters of UA were found to strongly inhibit expression of both MMP-2 and MMP-9 at micromolar concentrations in a DU-145 prostate tumor model ([Kondo et al. 2004](#)).

6.6.2. POLYPHENOLICS

Polyphenolic compounds are expected to play a key role in chemoprevention; thus, many in vitro cranberry studies focus on the berry's polyphenolic constituents. Cranberry extracts containing PACs and other flavonoids were reported to inhibit ODC activity linked with cell proliferation in mouse epithelial (ME-308) cells ([Kandil et al. 2002](#)). Characterization of an active subfraction revealed the presence of dimers and oligomers of catechin/epicatechin, monomeric catechins, and quercetin glycosides. Water-soluble cranberry polyphenolic extracts from commercial cranberry powder were found to inhibit proliferation of several human tumor cell lines ([Seeram et al. 2004](#)) including two oral (CAL27 and KB), four colon (HT-29, HCT-116, SW480, and SW620), and three prostate (RWPE-1, RWPE-2, and 22Rv1) cancer cell lines. Although both anthocyanin and PAC subfractions inhibited proliferation, the total polyphenolic extract was found to be more effective in this study. The primary polyphenolic constituents of cranberry are flavonols, anthocyanins, and PACs. These will be considered in [Sections 6.6.3](#) through [6.6.5](#).

6.6.3. QUERCETIN

Quercetin, a constituent of many fruits and vegetables, is widely reported to have antiproliferative and antineoplastic activities in vitro against a variety of cell lines ([Ranelletti, Maggiano, and Serra 2000](#); [Morrow et al. 2001](#); [Sesink, O'Leary, and Hollman 2001](#); [Lee, Huang, and Hwang 2002](#); [Harris et al. 2005](#); [Richter et al. 1999](#); [Volate et al. 2005](#)). Its

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

mechanisms of action include induction of apoptosis, with cell-cycle arrest in G₁ phase ([Richter, Ebermann, and Marian 1999](#); [Choi et al. 2001](#)); inhibition of epidermal growth factor (EGF) receptor expression and associated tyrosine kinase activity ([Richter, Ebermann, and Marian 1999](#); [Lee, Huang, and Hwang 2002](#)); reduction of Ras protein expression ([Ranelletti, Maggiano, and Serra 2000](#)); increased expression of endogenous inhibitors of MMPs ([Morrow et al. 2001](#)); and phytoestrogenic interaction with estrogen receptors α and β (ER α and ER β of human mammary MCF-7 cells; [Harris et al. 2005](#)). Quercetin is a constituent of an ethyl-acetate-soluble extract of cranberry fruit that inhibits tumor cell growth; it was nearly as effective as UA esters against the growth of MCF-7 breast adenocarcinoma, HT-29 colon adenocarcinoma, and K562 leukemia cell lines ([Murphy et al. 2003](#)). In a separate study, it was found to inhibit proliferation of HepG2 liver cancer cells ([He and Liu 2006](#)). Given the known properties of quercetin, it is a likely contributor to the observed in vitro anticancer activity of whole cranberry extracts. In vivo, quercetin glycosides are usually metabolized to sulfates or glucuronides ([Sesink, O'Leary, and Hollman 2001](#)). In a colon cancer study, a quercetin-enriched diet decreased formation of cancer precursor aberrant crypt foci fourfold in mice, and the evidence suggest that quercetin acted through induction of apoptosis via a mitochondrial pathway involving modulation of Bax and Bcl-2 protein expression ([Volate et al. 2005](#)).

6.6.4. ANTHOCYANINS

As powerful antioxidants, anthocyanins may be expected to inhibit oxidative processes linked with tumorigenesis. In vitro bioassays with cranberry anthocyanins show little direct antiproliferative or growth-inhibitory properties. Purified cyanidin-3-galactoside was evaluated in eight tumor cell lines in vitro using the sulforhodamine (SRB) assay and the highest concentration tested (250 $\mu\text{g/mL}$) showed less than 50% growth inhibition ([Murphy et al. 2003](#)). In another study, an anthocyanin subfraction from cranberry was found to limit growth in three prostate tumor lines (RWPE-1, RWPE-2, and 22Rv1) by 50–70%, but it did not significantly inhibit oral or colon tumor cell line proliferation ([Seeram et al. 2004](#)). Anthocyanins, including those from cranberry, have been implicated in the observed antiangiogenic properties of mixed berry extracts ([Roy et al. 2002](#); [Bagchi et al. 2004](#)). Anthocyanin-rich extracts from a mixture of berry fruits were reported to inhibit tumor necrosis factor α (TNF- α)-induced vascular endothelial growth factor (VEGF) expression and to decrease hemangioma formation and tumor growth ([Atalay et al. 2003](#)). Anthocyanins, though not especially cytotoxic, may nonetheless play a role in limiting carcinogenesis by inhibiting other activities related to tumor formation.

6.6.5. ANTICANCER ACTIVITIES OF CRANBERRY PROANTHOCYANIDINS

Many studies report cranberry PACs are major contributors to anticancer activity ([Kandil et al. 2002](#); [Ferguson et al. 2004](#); [Seeram et al. 2004](#)). In fractionation studies of whole fruit, a PAC fraction selectively inhibited the proliferation of H460 human large cell lung carcinoma, HT-29 colon adenocarcinoma, and K562 chronic myelogenous leukemia cells in vitro. A subfraction retaining the activity in those three cell lines was isolated and characterized by matrix-assisted laser desorption/ionization (MALDI)-TOF (time-of-flight)-MS, and was found to contain PAC oligomers composed primarily of four to seven epicatechin units with at least one or two A-type linkages between the units ([Neto et al. 2006](#)). In clonogenic assays for tumor colony formation with HT-29 and HCT-116 colon tumor cell lines, a dose-dependent decrease in the number and size of tumor colonies was observed when cells were treated with a PAC fraction prepared from

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

the early black variety of cranberry fruit ([Liberty et al. 2009](#)). The MALDI-TOF-MS characterization of this PAC fraction revealed that it was composed primarily of trimers through hexamers of epicatechin with both A- and B-type linkages. The fraction inhibited growth more effectively than the whole polyphenolic extract, with over 50% inhibition of colony formation in HCT116 observed at concentrations less than 10 µg/mL ([Liberty, Hart, and Neto 2007](#)).

The structures of PACs in cranberry fruit and cranberry products are complex, and further investigation is needed to determine any link between structure and activity of PAC. The MALDI-TOF-MS analysis found that PAC oligomer fractions from whole cranberry contained molecules up to 12 degrees of polymerization (DP) in size with as many as four A-type linkages. Although most fractions contained epicatechin units exclusively, some epigallocatechin unit masses were also detected ([Neto et al. 2006](#)). The effect of A-type linkages and oligomer size on the antitumor activity of PACs is the subject of ongoing studies. Earlier reports on PACs from various sources suggest that A-type linkages influence tumor-inhibitory and tumor-selectivity properties. A screening of small polyflavan-3-ols from different plants against GLC4 lung and COLO 320 colon carcinomas found that a trimer with an A-type linkage was more cytotoxic than dimers with A-type linkages and trimers with only B-type linkages ([Kolodziej et al. 1995](#)). In another study, wild blueberry PACs containing A-type linkages also showed selectivity, with a greater growth-inhibitory effect occurring with androgen-sensitive LNCaP prostate cancer cells than with androgen-insensitive DU145 cells ([Schmidt, Erdman, and Lila 2006](#)).

A recent study on cranberry PACs reports some promise for cranberry in alleviating cancer of the esophagus. A known risk factor for this type of cancer is acid reflux ([Herulf et al. 1999](#)). An in vitro model study employing SEG-1 human esophageal adenocarcinoma was used to investigate cranberry's effects on acid-induced cell proliferation. Cells were pretreated with PAC extract at concentrations of 25 or 50 µg/mL and then pulsed with acidified medium to simulate reflux. Maximum proliferation was induced 6 hours after treatment; further, proliferation was inhibited significantly in the PAC-treated cells. Annexin-V staining showed that apoptosis had occurred in treated cells, and flow cytometry experiments found these cells arrested at the G1 checkpoint ([Kresty, Howell, and Baird 2008](#)).

Recent findings suggest that cranberry PACs may work together with platinum drugs to limit the proliferation of ovarian cancer cells. In a study using a platinum-resistant SKOV-3 human ovarian adenocarcinoma cell line, treatment with a sublethal concentration of paraplatin together with an isolated cranberry PAC fraction was found to improve efficacy in decreasing cell proliferation as compared to treatment with PACs alone ([Singh et al. 2009](#)).

[Go to:](#)

6.7. POSSIBLE CHEMOPREVENTIVE MECHANISMS AND EFFECTS

Berry bioactives may act individually, additively, or synergistically to prevent cancer proliferation ([Seeram 2006](#)). Inhibition of tumorigenesis by cranberry can involve complementary or synergistic activities between the flavonols, PACs, UA, and anthocyanins, since all these compounds have been reported as showing antiproliferative properties. Possible mechanisms of action for chemoprevention supported by in vitro evidence include induction of apoptosis, decreased invasion of surrounding tissue due to MMP inhibition, reduction of ODC expression and activity, and inhibition of oxidative or inflammatory processes.

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

6.7.1. APOPTOSIS

Induction of apoptosis is thought to play a role in the tumor-inhibitory activity of dietary phytochemicals including resveratrol ([Joe et al. 2002](#)) and epigallocatechin gallate ([Yang et al. 1998](#)). Recent reports suggest that apoptosis may play a key role in cranberry's ability to limit tumor cell growth. This activity may be associated with the presence of quercetin, UA, and/or PACs, which are all known to induce apoptosis ([Baek et al. 1997](#); [Choi et al. 2001](#)). Dose-dependent induction of apoptosis by cranberry was observed in breast tumor models. An antiproliferative fraction from cranberry press cake also induced apoptosis in MDA-MB-435 breast tumor cells as determined by Annexin-V staining ([Ferguson et al. 2004](#)). An 80% aqueous acetone extract of whole cranberry fruit was reported to increase apoptosis in MCF-7 cells by 25% ([Sun and Liu 2006](#)), although at a concentration (50 mg/mL) much higher than would likely be encountered in vivo.

The whole polyphenolic extract of cranberry fruit was observed to increase apoptosis substantially in tumorigenic (MCF-7) breast cells as compared to nontumorigenic (MCF-10A) breast cells. At 250 µg/mL, the cranberry extract increased the baseline apoptosis rate to 92% in MCF-7 cells, without increasing apoptosis in MCF10A cells to a significant extent ([Griffin et al. 2005](#)). In colon tumor cell lines HCT116 and HT-29, treatment with UA or with cranberry PACs was found to induce significant rates of apoptosis at concentrations below 100 µg/mL ([Liberty, Hart, and Neto 2007](#)). Recent reports of the effects of cranberry PACs on human esophageal adenocarcinoma cells show that PACs induce apoptosis in this cell line ([Kresty, Howell, and Baird 2008](#)). Induction of apoptosis in SGC-7901 gastric cancer cells by an aqueous acetone extract of whole cranberry fruit was also reported ([Liu et al. 2009](#)). In a study of oral squamous cell carcinomas, a commercial cranberry powder reduced cell proliferation in SCC25 and CAL27 cell lines, and an upregulation of caspase-2 and caspase-8 messenger ribonucleic acid (mRNA) expression indicated that apoptosis played a role in reducing proliferation ([Chatelain et al. 2008](#)).

6.7.2. MATRIX METALLOPROTEINASE INHIBITION

Phytochemicals from whole cranberry fruit may act against cancers by limiting the processes involved in tumor invasion and metastasis, particularly the expression of MMPs involved in remodeling the extracellular matrix ([Pupa et al. 2002](#)). Both the whole cranberry polyphenolic extract and a cranberry PAC fraction were found to inhibit the expression of MMP-2 and MMP-9 in the DU145 prostate tumor cell line in a dose-dependent manner ([Neto et al. 2006](#)). Similar activity was reported for a flavonoid-rich extract of cranberry's close relative, the highbush blueberry (*V. angustifolium*; [Matchett et al. 2006](#)). Hydroxycinnamate esters of UA isolated from whole cranberry fruit were strong inhibitors of MMP-2 and MMP-9 protein expression, inhibiting expression significantly at concentrations of 10 µM or less ([Kondo et al. 2004](#)). This finding was consistent with the observed ability of UA to inhibit MMP expression in fibrosarcoma cells ([Cha et al. 1996](#)). Further studies are needed to determine the efficacy of cranberry and its PACs against tumor metastasis. Oligomers from grape seed have been found to possess antimetastasis activity both in vitro and in vivo ([Mantena, Baliga, and Katiyar 2006](#)).

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

6.7.3. MODULATION OF ORNITHINE DECARBOXYLASE

The biosynthesis and metabolism of polyamines (spermidine and spermine) involved in cell proliferation is controlled by enzymes such as ODC and spermidine/spermine N1-acetyltransferase, and ODC can be affected by dietary polyphenolics ([Singletary and Meline 2001](#)). Overexpression of these enzymes is observed in models of cancer where ODC can play a regulatory role in transformation, invasion, and angiogenesis ([Auvinen 1997](#)), and can be induced by proinflammatory agents such as lipopolysaccharides (LPSs) or tumor-promoters such as 12-*o*-tetradecanoyl phorbol-13-acetate (TPA). A cranberry fruit flavonoid fraction inhibits the activity of ODC in an ME-308 cell line, as determined by an assay measuring the conversion of substrate ([Kandil et al. 2002](#)). Cranberry was also found to influence the expression of ODC induced by LPS in an H-ras transformed mouse fibroblast model ([Matchett et al. 2005](#)). A whole cranberry polyphenolic extract produced a dose-dependent inhibition of LPS-induced ODC expression, and induction by LPS was abolished at extract concentrations of 100 µg/mL or less ([Matchett et al. 2005](#)).

6.7.4. HELICOBACTER PYLORI INHIBITION

H. pylori infection is positively associated with the incidence of gastric cancer ([Uemura et al. 2001](#)); thus, the prevention of this infection may reduce cancer risk. Antibacterial adhesion studies demonstrate that in addition to inhibiting *Escherichia coli* adhesion, cranberry components inhibit adhesion of *H. pylori* to human gastric mucus ([Burger et al. 2000](#)). A randomized, double-blind, placebo-controlled trial provided some clinical support for this finding, with significantly lower levels of *H. pylori* infection observed in adults consuming cranberry juice ([Zhang et al. 2005](#)). The cranberry polyphenolic extract and other polyphenol-rich juices had a bacteriostatic effect on the growth of *H. pylori* in vitro, with morphological changes in bacteria seen at concentrations of 1 mg/mL ([Matsushima et al. 2008](#)).

6.7.5. EVIDENCE FROM ANIMAL MODELS

Tumor growth inhibition by cranberry in an animal model was first reported in 2006 ([Ferguson et al. 2006](#)) by a group of Canadian researchers. In this study, explants of U87 glioblastoma, HT-29 colon carcinoma, or DU145 prostate carcinoma were established in female Balb/c mice. Mice in the treatment groups were intraperitoneally injected with either a flavonoid-rich aqueous extract from cranberry press cake or a PAC fraction prepared from the whole fruit extract. Dosages of 100 mg/kg body weight PAC fraction or 250 mg/kg press cake extract were administered 10 times over a period of 24 days. Both treatments resulted in up to 40% reduction in the time required for U87 glioblastoma tumors to reach milestone sizes. Flow cytometry experiments showed that the extracts arrest U87 cells in G1 phase after 24 hours, reducing the number of cells continuing to S phase. In mice given HT-29 tumor explants, the PAC treatment group exhibited significantly reduced tumor volume over the first 40 days compared to control. Proanthocyanidin treatment was also found to slow the growth of tumors in the DU145 group and induce complete regression of these tumors in the two treatment groups of mice ([Ferguson et al. 2006](#)).

Several animal studies have since appeared that examine the effects of cranberry treatment on models of cancer. A 2008 study used immune-competent syngeneic mice to investigate the

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

effects of a nondialyzable (NDM) high-molecular-weight fraction from cranberry juice on the development of lymphoma ([Hochman et al. 2008](#)). The fraction was presumed to contain polyphenolic oligomers in the 12–30K weight range; therefore, exact structures could not be determined. Balb/c female mice were inoculated with Rev-2-T-6 lymphoma cells. The cranberry NDM fraction was injected into mice at nontoxic doses (2 or 4 mg) every 2 days for 2 weeks. At the end of the experiment, cranberry-treated mice showed no tumor development in comparison with the control group, in which 80% developed tumors within 3 weeks. The treated mice also showed an increase in the production of antibodies against lymphoma cells. In another study, Balb/c nu/nu mice were injected with human gastric cancer cell line SGC-7901 that had been pretreated with an acetone-soluble cranberry extract at doses ranging from 5 to 40 mg. After 4 weeks, the control group developed tumors (xenografts), whereas no tumors were observed in the highest cranberry dosage group and tumor size was reduced in mice receiving cranberry extract doses as low as 10 mg ([Liu et al. 2009](#)).

In a bladder cancer study that used Fischer-344 rats as a model, rats were treated with N-butyl-N(4-hydroxybutyl)nitrosamine to induce cancer. Rats received cranberry juice concentrate via gavage at doses of 1.0 or 0.5 mL/rat/day (nontoxic doses) for up to 6 months. Both treatment groups showed a significant decrease in the number of bladder lesions, particularly papillomas, and the higher dosage group showed a significant decrease in number of carcinomas and a 31% decrease in the total weight of bladder lesions ([Prasain et al. 2008](#)). Quercetin, a key constituent of cranberry juice, was not detected in the plasma; however, both quercetin and its metabolite methyl quercetin were detected in the urines of the treated mice.

6.7.6. ANTI-INFLAMMATORY ACTIVITIES

The anti-inflammatory properties of phytochemicals may have an impact on many diseases including certain cancers. Cyclooxygenase (COX) is a key enzyme in the biosynthetic pathway to prostaglandins, which have many physiological roles, including the production of an inflammatory response. The COX-1 isozyme is expressed constitutively in all cells, whereas expression of COX-2 can be induced in response to inflammatory stimuli. The COX-2 is highly expressed in tumor tissues ([Bottone et al. 2004](#)), and studies show that nonsteroidal anti-inflammatory drugs (NSAIDs), for example, Sulindac, have a chemopreventive effect against colon cancer in cellular and animal models ([Sheng et al. 1997](#); [Fournier and Gordon 2000](#); [Bottone et al. 2004](#)). As COX-2 overexpression is thought to play a role in promoting certain cancers, inhibition of COX-2 activity or expression presents another potential route to chemoprevention. Several individual constituents of cranberry may have anti-inflammatory properties. Inhibition of COX activity by anthocyanins, including those found in cranberry, was reported ([Seeram et al. 2001](#)). Pure cyanidin is an effective COX-2 inhibitor, which reduces activity by 47%, with activity superior to other anthocyanins or catechins ([Seeram, Zhang, and Nair 2003](#)). The observed COX-1 inhibition may also be relevant to cancer, since some evidence suggests that COX-1 specific inhibitors are as effective as COX-2 specific inhibitors in decreasing events related to tumor development ([Bottone et al. 2004](#)).

The question of whether cranberry can decrease the expression of COX-1 or COX-2 in cellular models remains to be answered; to date, no published studies are available that evaluate the effects of cranberry on COX expression in cancer models. Inhibition of COX-2 expression, if observed, could contribute to anticancer activity. Considering the known effects of compounds

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

found in cranberry fruit, modulation of COX-2 expression and associated pathways is likely to be beneficial. Both UA and quercetin are established inhibitors of cellular COX expression ([Subbaramaiah et al. 2000](#); [O'Leary et al. 2004](#)). The anti-inflammatory actions of triterpenes including UA have been reviewed ([Safayhi and Sailer 1997](#)), and studies support an anti-inflammatory role both in vitro ([Ringbom et al. 1998](#); [Subbaramaiah et al. 2000](#)) and in vivo ([Recio et al. 1995](#); [Baricevic et al. 2001](#)). UA inhibited COX-2 transcription in a human mammary oncogenic epithelial cell line (184B5/HER) by a mechanism believed to involve the protein kinase C signal transduction pathway ([Subbaramaiah et al. 2000](#)). Several anti-inflammatory activities of quercetin have also been reported. Quercetin reduced COX-2 mRNA expression in Caco-2 colon cancer cells, and both quercetin and its metabolite quercetin-3'-sulfate inhibited COX-2 activity ([O'Leary et al. 2004](#)). In a rat model of colitis, quercetin delivered in the form of rutin inhibited the TNF- α -dependent activation of nuclear factor κ B (NF- κ B), a transcription factor involved in the control of cell proliferation and inflammation, in a dose-dependent manner ([Kim et al. 2005](#)). Similarly, UA has been reported to suppress NF- κ B activation ([Shishodia et al. 2003](#)). Quercetin has been reported to inhibit expression of inducible nitric oxide synthase, a promoter of inflammation that has also been linked to tumor angiogenesis, in cellular models ([García-Mediavilla et al. 2007](#)).

[Go to:](#)

6.8. CRANBERRIES AND URINARY HEALTH

The use of cranberry juice to prevent UTIs has a long history that was for many years supported mainly by anecdotal evidence. This is no longer the case: A body of scientific evidence has accumulated to support the use of cranberry in the maintenance of urinary tract health. Studies started appearing in the 1980s demonstrating the ability of cranberry juice to prevent adherence of *E. coli* bacteria to uroepithelial cells and other eukaryotic cells ([Sobota 1984](#); [Zafriri et al. 1989](#); [Ofek et al. 1991](#)). As type 1-fimbriated bacteria were susceptible to the fructose in citrus fruit juices as well, the effect on type P-fimbriated *E. coli* was observed to be specific to cranberry ([Zafriri et al. 1989](#)) and other *Vaccinium*. During the mid-1990s, a clinical study conducted by [Avorn et al. \(1994\)](#) on the female residents of a long-term care facility found a significant decrease in bacteria in the urine after 1 month of cranberry juice consumption. Since then, at least 15 clinical trials have evaluated the prophylactic effects of cranberry against urinary infections in a variety of populations. These studies are the subject of several detailed review articles ([Howell 2002](#); [Jepson and Craig 2007](#); [Guay 2009](#)).

For many years, scientists and health practitioners believed that the antibacterial effects of cranberry juice were due to acidification of urine by hippuric acid produced by the metabolism of the quinic acid in cranberries ([Blatherwick 1914](#); [Bodel, Cotran, and Kass 1959](#)). However, this claim was never substantiated. Studies correlating urinary pH with cranberry juice consumption show either no significant change in urine acidity or only a slight reduction in pH, which is insufficient to cause a bacteriostatic effect ([Howell 2002](#)). Researchers began to examine other possible mechanisms of action, which led to the discovery of bacterial antiadhesion properties ([Sobota 1984](#)). Studies on antiadherence of uropathogenic P-fimbriated *E. coli* (UPEC) responsible for the majority of UTIs found that a high-molecular-weight NDM from cranberry inhibited adhesion ([Ofek et al. 1991](#)). Bioassay-guided fractionation of cranberry targeting the active compounds found that cranberry tannins ([Howell](#)

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

[et al. 1998](#)) blocked adherence of P-fimbriated *E. coli* to uroepithelial cells. The structures of these compounds were determined by nuclear magnetic resonance (NMR) analysis to be polyflavan-3-ol or PAC trimers composed of epicatechin units with an A-type linkage ([Foo et al. 2000](#)). The A-type linkage between monomer units ([Figure 6.3](#)), which features two linkage sites between the units ($4\beta\rightarrow 8$ and $2\beta\rightarrow O\rightarrow 7$ interflavanoid bonds), is a structural feature common to PACs from *Vaccinium* fruit. Proanthocyanidins from most other sources including cocoa and grape seeds contain primarily B-type ($4\beta\rightarrow 8$) linkages ([Neto 2007](#)), as shown in [Figure 6.3](#). Cranberry PACs are primarily dimers, trimers, and larger oligomers of epicatechin units containing both A- and B-type linkages; therefore, their three-dimensional structures are diverse. Although cranberry NDM has been cited more in recent studies for its ability to inhibit cellular adhesion of *H. pylori* ([Burger et al. 2000](#)) and the development of lymphoma ([Hochman et al. 2008](#)), no information is available on the molecular structures of its components.

The A-type linkage may hold the key to the antiadhesion activity of cranberry PACs. A comparison study of PACs isolated from several food sources, including cranberry, apple, grape, green tea, and chocolate, found that cranberry PACs prevented *E. coli* adhesion at the lowest concentration tested (60 $\mu\text{g/mL}$). Among the others, grape PACs showed antiadherence properties, but only at a much higher dose (1200 $\mu\text{g/mL}$), and the other food sources showed no activity ([Howell et al. 2005](#)). This study further found antiadhesion activity in human urine following consumption of cranberry juice cocktail, but not after consumption of other PAC sources. A randomized, double-blind, placebo-controlled crossover trial with 20 healthy volunteers was conducted to determine whether urine collected after cranberry consumption inhibited UPEC adherence to uroepithelial bladder cells. Subjects received a single dose of cranberry juice (750 or 250 mL mixed with 500 mL water) or placebo at night, and urine was collected in the morning and screened for antiadhesion against six UPEC strains. A significant and dose-dependent decrease in adherence of bacteria was observed in the urine of subjects who consumed cranberry ([Di Martino et al. 2006](#)).

Several studies have examined the phenomenon of antibacterial adhesion by cranberry constituents in an attempt to better understand what happens at a submicroscopic level. Atomic force microscopy was used to measure the effect of cranberry juice exposure on bacterial surface characteristics and adhesion forces in a P-fimbriae-expressing *E. coli*(HB101pDC1) and a nonfimbriated strain. A decrease in fimbrial length was measured after a short exposure to cranberry juice, with a greater biopolymer density recorded near the cell wall ([Liu et al. 2006](#)). Adhesion forces decreased in proportion to cranberry juice concentration. No significant effect of cranberry on surface polymers or adhesive forces was observed with nonfimbriated *E. coli*. Later studies by this group showed that the effects of culturing the bacteria in cranberry juice or PAC extract on bacterial adhesion forces were reversible ([Pinzon-Arango, Liu, and Camesano 2009](#)). Interestingly, a more marked decrease in adhesive forces was observed with the juice cocktail than with the PAC fraction, suggesting that other components in the juice play a role in reducing bacterial adhesion. Another study found that morphological changes occurred when *E. coli* bacteria were grown in the presence of cranberry juice or PAC extract ([Johnson et al. 2008](#)). The authors observed a decrease in visible P-fimbriae and downregulation of gene expression associated with flagellar basal body rod and motor proteins.

Clinical studies have demonstrated the efficacy of consuming cranberry juice or solids in UTI prevention for various populations, including women with recurrent UTIs ([Walker et al.](#)

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

[1997](#); [Stothers 2002](#); [Bailey et al. 2007](#)), pregnant women, ([Wing et al. 2008](#)), the elderly ([Avorn et al. 1994](#); [McMurdo et al. 2005](#); [McMurdo et al. 2009](#)), and children ([Ferrara et al. 2009](#)). A randomized, double-blind, placebo-controlled study of women aged 28–44 years with recurring UTIs using 400 mg of cranberry solids or placebo for 3 months found that 70% of the subjects had fewer UTIs while on cranberry ([Walker et al. 1997](#)). In a study of women aged 25–70 years with a history of high UTI recurrence (six or more in the previous year), it was found that consumption of 200 mg of concentrated cranberry extract standardized to 30% phenolics twice a day for 12 weeks prevented UTI recurrence in all subjects for the duration of the study ([Bailey et al. 2007](#)). A follow-up study of these subjects 2 years later found that those who continued to take cranberry remained free of infection. In pregnant women, for which asymptomatic bacteriuria can cause adverse perinatal outcomes if not detected and treated, consumption of 27% cranberry juice cocktail (240 mL, thrice daily) resulted in 57% reduction in asymptomatic bacteriuria and 41% reduction in UTI ([Wing et al. 2008](#)).

In addition to Avorn's landmark study on elderly women ([Avorn et al. 1994](#)), a randomized, double-blind, placebo-controlled study of hospitalized patients over 60 years of age found that the group consuming 25% cranberry juice cocktail (150 mL twice daily for 35 days or until discharge) had approximately half the occurrence of symptomatic UTI ([McMurdo et al. 2005](#)). A follow-up study by this group comparing treatment with 500 mg of cranberry extract to that with low doses of the antibiotic trimethoprim in older women found that both treatments reduced recurrent UTI. Trimethoprim treatment had only a limited advantage in this regard, and the cranberry group reported less adverse effects ([McMurdo et al. 2009](#)). Further clinical studies in such populations continue to provide information for health practitioners on the most effective forms and dosing regimens of cranberry against UTI.

[Go to:](#)

6.9. FUTURE DIRECTIONS FOR RESEARCH ON CRANBERRIES AND DISEASE

Evidence from in vitro, in vivo, and clinical studies suggests that cranberry and its phytochemicals may have a mitigating effect on UTIs, cardiovascular diseases, and cancer. Mechanisms of action of cranberry range from antioxidative and anti-inflammatory actions to induction of cellular apoptosis; modulation of protein synthesis and gene expression involved in cell proliferation; prevention of bacterial adhesion and formation of biofilms that lead to infection; and effects on plasma lipoprotein levels, antioxidant status, and glucose metabolism. Future research should continue to examine cranberry's role in regulating these processes and address how the unique blend of phytochemicals found in cranberry fruit and juice works together. Cranberry's efficacy depends largely on the bioavailability of its phytochemicals to various tissues, which is another topic for further research.

[Go to:](#)

ACKNOWLEDGMENTS

Catherine Neto would like to acknowledge the assistance of University of Massachusetts, Dartmouth (UMD) science librarian Elizabeth Winiarz in gathering materials for this article, as well as the support of the UMD Cranberry Research Program and the Cranberry Institute. Joe

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

Vinson would like to thank Debra Jennings for secretarial assistance, Yvette Vinson for proofreading, and the Cranberry Institute for financial support.

REFERENCES

1. Aherne S.A, O'Brien N.M. Dietary flavonols: Chemistry, food content, and metabolism. *Nutr.* 2002;18:75–81. [\[PubMed\]](#)
2. Andersson D, Liu J.J, Nilsson A, Duan R.D. Ursolic acid inhibits proliferation and stimulates apoptosis in HT29 cells following activation of alkaline sphingomyelinase. *Anticancer Res.* 2003;23:3317–22. [\[PubMed\]](#)
3. Atalay M, Gordillo G, Roy S, editors. Anti-angiogenic property of edible berry in a model of hemangioma. *FEBS Lett.* 2003;544:252–7. [\[PubMed\]](#)
4. Auvinen M. Cell transformation, invasion, and angiogenesis: A regulatory role for ornithine decarboxylase and polyamines? *J Nat Cancer Inst.* 1997;89:533–7. [\[PubMed\]](#)
5. Avorn J, Monane M, Gurwitz J.H, Glynn R.J, Choodnovskiy I, Lipsitz L.A. Reduction of bacteriuria and pyuria after ingestion of cranberry juice. *J Am Med Assoc.* 1994;271:751–4. [\[PubMed\]](#)
6. Baek J.H, Lue Y.S, Kang C.M, editors. Intracellular Ca²⁺ release mediates ursolic acid-induced apoptosis in human leukemic HL-60 cells. *Int J Cancer.* 1997;73:725–8. [\[PubMed\]](#)
7. Bagchi D, Sen C.K, Bagchi M, Atalay M. Anti-angiogenic, antioxidant and anticarcinogenic properties of a novel anthocyanin-rich berry extract formula. *Biochem Trans Biokhimiya.* 2004;69:79–80. [\[PubMed\]](#)
8. Bailey D.T, Dalton C, Daugherty F.J, Tempesta M.S. Can a concentrated cranberry extract prevent recurrent urinary tract infections in women? A pilot study. *Phytomed.* 2007;14:237–41. [\[PubMed\]](#)
9. Baricevic D, Sosa S, Della Loggia R, editors. Topical antiinflammatory activity of *Salvia officinalis* L. leaves: The relevance of ursolic acid. *J Ethnopharmacol.* 2001;75:125–32. [\[PubMed\]](#)
10. Baur J.A, Sinclair D.A. Therapeutic potential of resveratrol: The in vivo evidence. *Nat Rev Drug Discov.* 2006;5:493–506. [\[PubMed\]](#)
11. Bilyk A, Sapers G.M. Varietal differences in the quercetin, kaempferol, and myricetin contents of highbush blueberry, cranberry, and thornless blackberry fruits. *J Agric Food Chem.* 1986;34:585–8.
12. Bitsch I, Janssen M, Netzel M, Strass G, Frank T. Bioavailability of anthocyanidin-3-glycosides following consumption of elderberry extract and black currant juice. *Int J Clin Pharmacol Ther.* 2004;42:293–300. [\[PubMed\]](#)
13. Blatherwick N.R. The specific role of foods in relation to the composition of the urine. *Arch Intern Med.* 1914;14:409–50.
14. Bodel P.T, Cotran R, Kass E.H. Cranberry juice and the antibacterial action of hippuric acid. *J Lab Clin Med.* 1959;54:881–8. [\[PubMed\]](#)
15. Bomser J, Madhavi D.L, Singletary K, Smith M.A. In vitro anticancer activity of fruit extracts from *Vaccinium* species. *Planta Med.* 1996;62:212–6. [\[PubMed\]](#)

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

16. Bottone F.G, Martinez J.M, Alston-Mills B, Eling T.E. Gene modulation by Cox-1 and Cox-2 specific inhibitors in human colorectal carcinoma cancer cells. *Carcinogenesis*. 2004;25:349–57. [[PubMed](#)]
17. Burger O, Ofek I, Tabak M, Weiss E.I, Sharon N, Neeman I. A high molecular mass constituent of cranberry juice inhibits *Helicobacter pylori* adhesion to human gastric mucus. *FEMS Immunol Med Microbiol*. 2000;29:295–301. [[PubMed](#)]
18. Cape Cod Cranberry Grower's Association. "History of Cranberries." <http://www.cranberries.org/cranberries/history.html> (accessed October 5, 2009).
19. Celik H, Özgen M, Serçe S, Kaya C. Phytochemical accumulation and antioxidant capacity at four maturity stages of cranberry fruit. *Sci Hortic*. 2008;117:345–8.
20. Cha H.J, Bae S.K, Lee H.Y, editors. Anti-invasive activity of ursolic acid correlates with the reduced expression of matrix metalloproteinase-9 (MMP-9) in HT1080 human fibrosarcoma cells. *Cancer Res*. 1996;56:2281–4. [[PubMed](#)]
21. Chambers B.K, Camire M.E. Can cranberry supplementation benefit adults with type 2 diabetes? *Diabetes Care*. 2003;26:2695–6. [[PubMed](#)]
22. Chatelain K, Phippen S, McCabe J, Teeters C.A, O'Malley S, Kingsley K. Evid Based Complement Altern Med. 2008. Cranberry and grape seed extracts inhibit the proliferative phenotype of oral squamous cell carcinomas. Advance access, published on July 23, 2008; doi:10.1093/ecam/nen047.
23. Choi Y.H, Baek J.H, Yoo M.A, Chung H.Y, Kim N.D, Kim K.W. Induction of apoptosis by ursolic acid through activation of caspases and down-regulation of c-IAPs in human prostate epithelial cells. *Int J Oncol*. 2000;17:565–71. [[PubMed](#)]
24. Choi J, Kim J, Lee J, editors. Induction of cell cycle arrest and apoptosis in human breast cancer cells by quercetin. *Int J Oncol*. 2001;19:837–44. [[PubMed](#)]
25. Croteau R.J, Fagerson I.S. Major volatile components of the juice of American cultivar and storage temperature. *J Agric Food Chem*. 1968;49:969–74.
26. Cunningham D.G, Vnnozzi S.A, Turk R, Roderick R, O'Shea E, Brilliant K. Cranberry phytochemicals and their health benefits nutraceutical beverages. Washington, DC: American Chemical Society; ACS Symposium Series. 2003;vol. 871:35–51. Chap. 4.
27. Di Martino P, Agniel R, David K, Templar C, Gaillard J.L, Denys P, Botto H. Reduction of *Escherichia coli* adherence to uroepithelial bladder cells after consumption of cranberry juice: A double-blind, randomized placebo-controlled cross-over trial. *World J Urol*. 2006;24:21–7. [[PubMed](#)]
28. Duthie S.J, Jenkinson A.M, Crozier A, editors. The effects of cranberry juice consumption on antioxidant status and biomarkers relating to heart disease and cancer in healthy human volunteers. *Eur J Nutr*. 2006;45:113–22. [[PubMed](#)]
29. Duthie G.G, Kyle J.A, Jenkinson A.M, Duthie S.J, Baxter G.J, Paterson J.R. Increased salicylate concentrations in urine of human volunteers after consumption of cranberry juice. *J Agric Food Chem*. 2005;53:2897–9000. [[PubMed](#)]
30. Ellingsen I, Hjerkin E.M, Seljeflot I, Arnesen H, Tonstad S. Consumption of fruit and berries is inversely associated with carotid atherosclerosis in elderly men. *Br J Nutr*. 2008;99:674–81. [[PubMed](#)]
31. Es-Saady D, Simon A, Jayat-Vignoles C, Chulia A.J, Delage C. MCF-7 cell cycle arrested at G1 through ursolic acid and increased reduction of tetrazolium salts. *Anticancer Res*. 1996;16:481–6. [[PubMed](#)]

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

32. Ferguson P, Kurowska E, Freeman D.J, Chambers A.F, Koropatnick D.J. A flavonoid fraction from cranberry extract inhibits proliferation of human tumor cell lines. *J Nutr.* 2004;134:1529–35. [[PubMed](#)]
33. Ferguson P.J, Kurowska E.M, Freeman D.J, Chambers A.F, Koropatnick D.J. In vivo inhibition of growth of human tumor lines by flavonoid fractions from cranberry extract. *Nutr Cancer.* 2006;56:86–94. [[PubMed](#)]
34. Ferrara P, Romaniello L, Vitelli O, Gatto A, Serva M, Cataldi L. Cranberry juice for the prevention of recurrent urinary tract infections: A randomized controlled trial in children. *Scand J Urol Nephrol.* 2009;43:369–372. [[PubMed](#)]
35. Foo L.Y, Lu Y, Howell A.B, Vorsa N. A-type proanthocyanidin trimers from cranberry that inhibit adherence of uropathogenic P-fimbriated. *Escherichia coli.* *J Nat Prod.* 2000a;63:1225–8. [[PubMed](#)]
36. Foo L.Y, Lu Y, Howell A.B, Vorsa N. The structure of cranberry anthocyanidins proanthocyanidins which inhibit adherence of uropathogenic P-fimbriated *Escherichia coli* in vitro. *Phytochemistry.* 2000b;54:173–81. [[PubMed](#)]
37. “Cranberries: Cranberry Trivia”. Foodreference.com. <http://www.foodreference.com/html/fcranberries.html> (accessed October 5, 2009).
38. “Cranberries”. FoodReference.com. <http://www.foodreference.com/html/fcranberries.html> (accessed October 5 2009)
39. Fournier D.B, Gordon G.B. COX-2 and colon cancer: Potential targets for chemoprevention. *J Cell Biochem.* 2000;77:97–102. [[PubMed](#)]
40. Frank T, Netzel M, Strass G, Bitsch R, Bitsch I. Bioavailability of anthocyanidin-3-glucosides following consumption of red wine and red grape juice. *Can J Physiol Pharmacol.* 2003;81:423–35. [[PubMed](#)]
41. Freshfel Europe. Freshfel Fruit and Vegetable Production, Trade, Supply and Consumption Monitor in the EU-27. 2009. Brussels, Belgium, Freshfel Europe.
42. Fuleki T, Francis F.J. Quantitative methods for anthocyanins: Purification of cranberry anthocyanins. *J Food Sci.* 1968;33:266–9.
43. García-Mediavilla V, Crespo I, Collado P.S, Esteller A, Sánchez-Campos S, Tunòn M.J, González-Gallego J. The anti-inflammatory flavones quercetin and kaempferol cause inhibition of inducible nitric oxide synthase, cyclooxygenase-2 and reactive C-protein, and down-regulation of the nuclear factor kappa B pathway in Chang liver cells. *Eur J Pharmacol.* 2007;557:221–9. [[PubMed](#)]
44. Giordano L, Coletta W, Rapisard P, Donati M.B, Rotilio D. Development and validation of an LC-MS/MS analysis for simultaneous determination of delphinidin-3-glucoside, cyanidin-3-glucoside and cyanidin-3-(6-malonylglucoside) in human plasma and urine after blood orange juice administration. *J Sep Sci.* 2007;30:3127–36. [[PubMed](#)]
45. Griffin L, Rego S, Correiro E, Neto C, Hart P. San Diego, CA: American Society for Cell Biology; Induction of Apoptosis in Tumor Cell Lines by Polyphenolic Compounds Isolated from *Vaccinium Macrocarpon*. 2005
46. Gu L, Kelm M.A, Hammerstone J.F, editors. Concentrations of proanthocyanidins in common foods and estimations of normal consumption. *J Nutr.* 2004;134:613–7. [[PubMed](#)]
47. Gu L, Kelm M.A, Hammerstone J.F, Beecher G, Holden J, Haytowitz D, Prior R.L. Screening of foods containing proanthocyanidins and their structural characterization

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

- using LC-MS/MS and thiolytic degradation. *J Agric Food Chem.* 2003;51:7513–21. [[PubMed](#)]
48. Guay D.R. Cranberry and urinary tract infections. *Drugs.* 2009;69:775–807. [[PubMed](#)]
49. Guthrie N. Effect of cranberry juice and products on human breast cancer cell growth. *Experimental Biology.* 2003 San Diego, CA.
50. Häkkinen S.H, Kärenlampi S.O, Heinonen I.M, Mykkänen H.M, Törrönen A.R. Content of the flavonols quercetin, myricetin, and kaempferol in 25 edible berries. *J Agric Food Chem.* 1999;47:2274–9. [[PubMed](#)]
51. Harnly J.M, Doherty R.F, Beecher G.R, Holden J.M, Haytowitz D.B, Bhagwat S, Gebhardt S. Flavonoid content of U.S. fruits, vegetables, and nuts. *J Agric Food Chem.* 2006;54:9966–77. [[PubMed](#)]
52. Harris D.M, Besselink E, Henning S.M, Go V. L. W, Heber D. Phytoestrogens induce differential estrogen receptor alpha- or beta-mediated responses in transfected breast cancer cells. *Exp Biol Med.* 2005;230:558–68. [[PubMed](#)]
53. He X, Liu R.H. Cranberry phytochemicals: Isolation, structure elucidation and their antiproliferative and antioxidant activities. *J Agric Food Chem.* 2006;54:7069–74. [[PubMed](#)]
54. Herulf M, Lagergren A, Ljung T, Morcos E, Wiklund N.P, Lundberg J.O, Weitzberg E. Increased nitric oxide in infective gastroenteritis. *J Infect Dis.* 1999;180:542–5. [[PubMed](#)]
55. Hochman N, Hourri-Haddad Y, Koblinski J, editors. Cranberry juice constituents impair lymphoma growth and augment the generation of antilymphoma antibodies in syngeneic mice. *Nutr Cancer.* 2008;60:511–7. [[PubMed](#)]
56. Howell A. Cranberry proanthocyanidins and the maintenance of urinary tract health. *Crit Rev Food Sci Nutr.* 2002;42S:273–8. [[PubMed](#)]
57. Howell A.B, Reed J.D, Krueger C.G, Winterbottom R, Cunningham D.G, Leahy M. A-type cran-berry proanthocyanidins and uropathogenic bacterial anti-adhesion activity. *Phytochem.* 2005;66:2281–91. [[PubMed](#)]
58. Howell A.B, Vorsa N, Der Marderosian A, Foo L.Y. Inhibition of the adherence of P-fimbriated *Escherichia coli* to uroepithelial-cell surfaces by proanthocyanidin extracts from cranberries. *N Eng J Med.* 1998;339:1085–6. [[PubMed](#)]
59. Jensen H.D, Krogfelt K.A, Cornett C, Hansen S.H, Christensen S.B. Hydrophilic carboxylic acids and iridoid glycosides in the juice of American and European cranberries (*Vaccinium macrocarpon* and *V oxycoccos*), lingonberries (*V. vitis-idaea*), and blueberries (*V. myrtillus*). *J Agric Food Chem.* 2002;50:6871–4. [[PubMed](#)]
60. Jepson R.G, Craig J.C. A systematic review of the evidence for cranberries and blueberries in UTI prevention. *Mol Nutr Food Res.* 2007;51:738–45. [[PubMed](#)]
61. Joe A.K, Liu H, Suzui M, Vural M.E, Xiao D, Weinstein I.B. Resveratrol induces growth inhibition, S-phase arrest, apoptosis, and changes in biomarker expression in several human cancer cell lines. *Clin Cancer Res.* 2002;8:893–903. [[PubMed](#)]
62. Johnson B.J, Lin B, Dinderman M.A, Rubin R.A, Malanoski A.P, Ligler F.S. Impact of cranberry on *Escherichia coli* cellular surface characteristics. *Biochem Biophys Res Comm.* 2008;377:992–4. [[PubMed](#)]
63. Kähkönen M.P, Hopia A.I, Heinonen M. Berry phenolics and their antioxidant activity. *J Agric Food Chem.* 2001;49:4076–82. [[PubMed](#)]

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

64. Kandil F.E, Smith M. A. L, Rogers R.B, editors. Composition of a chemopreventive proantho-cyanidin-rich fraction from cranberry fruits responsible for the inhibition of TPA-induced ODC activity. *J Agric Food Chem.* 2002;50:1063–9. [[PubMed](#)]
65. Kim D.K, Baek J.H, Kang C.M, editors. Apoptotic activity of ursolic acid may correlated with the inhibition of initiation of DNA replication. *Int J Cancer.* 2000;87:629–36. [[PubMed](#)]
66. Kim H, Kong H, Choi B, editors. Metabolic and pharmacological properties of rutin, a dietary quercetin glycoside, for treatment of inflammatory bowel disease. *Pharm Res.* 2005;22:1499–509. [[PubMed](#)]
67. Kim K.A, Lee J.S, Park H.J, editors. Inhibition of cytochrome P450 activities by oleanolic acid and ursolic acid in human liver microsomes. *Life Sci.* 2004;74:2769–79. [[PubMed](#)]
68. Kolodziej H, Haberland C, Woerdenbag H.J, Konings A. W. T. Moderate cytotoxicity of proanthocyanidins to human tumour cell lines. *PhytotherRes.* 1995;9:410–5.
69. Kondo M. *Phytochemical Studies of Extracts from Cranberry (Vaccinium macrocarpon) with Anticancer, Antifungal and Cardioprotective Properties.* 2006 North Dartmouth, MA: University of Massachusetts Dartmouth.
70. Kondo M, Lamoureaux T.L, Neto C.C. *J Nutr.* 2004. Proanthocyanidins, anthocyanins and triterpenoids from cranberry fruits: Antitumor activity and effects on matrix metalloproteinase expression. Abstract. 12S:3538S.
71. Kresty L.A, Howell A.B, Baird M. Cranberry proanthocyanidins induce apoptosis and inhibit acid-induced proliferation of human esophageal adenocarcinoma cells. *J Agric Food Chem.* 2008;56:676–80. [[PubMed](#)]
72. Lee I.T, Chan Y.C, Lin C.W, Lee W.J, Sheu W.H. Effect of cranberry extracts on lipid profiles in subjects with type 2 diabetes. *Diabet Med.* 2008;25:1473–7. [[PubMed](#)]
73. Lee L.T, Huang Y.T, Hwang J.J. Blockade of the epidermal growth factor receptor tyrosine kinase activity by quercetin and luteolin leads to growth inhibition and apoptosis of pancreatic tumor cells. *Anticancer Res.* 2002;22:1615–27. [[PubMed](#)]
74. Lee I, Lee J, Lee Y.H, Leonard J. Ursolic-acid induced changes in tumorgrowth, O2 consumption, and tumor interstitial fluid pressure. *Anticancer Res.* 2001;21:2827–33. [[PubMed](#)]
75. Liberty A.M, Hart P.E, Neto C.C. Ursolic acid and proanthocyanidins from cranberry (*Vaccinium macrocarpon*) inhibit colony formation and proliferation in HCT-116 and HT-29 colon and MCF-7 breast tumor cells. 233rd National Meeting of the American Chemical Society, Chicago, IL. 2007
76. Liberty A.M, Amoroso J.W, Hart P.E, Neto C.C. Cranberry PACs and triterpenoids: anti-cancer activities in colon tumor cell lines. *Proceedings of the Second International Symposium on Human Health Effects of Fruits and Vegetables. Acta Horticulturae.* 2009;841:61–66.
77. Liu Y, Black M.A, Caron L, Camesano T.A. Role of cranberry juice on molecular-scale surface characteristics and adhesion behavior of *Escherichia coli*. *Biotechnol Bioeng.* 2006;93:297–305. [[PubMed](#)]
78. Liu M.L, Lin L.Q, Song B.B, editors. Cranberry phytochemical extract inhibits SGC-7901 cell growth and human tumor xenografts in Balb/c nu/nu mice. *J Agric Food Chem.* 2009;57:762–8. [[PubMed](#)]

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

79. Manach C, Scalbert A, Morand C, Rèmesy C, Jimenez L. Polyphenols: Food sources and bioavailability. *Am J Clin Nutr.* 2004;79:727–47. [[PubMed](#)]
80. Mantena S.K, Baliga M.S, Katiyar S.K. Grape seed proanthocyanidins induce apoptosis and inhibit metastasis of highly metastatic breast carcinoma cells. *Carcinogenesis.* 2006;27:1682–91. [[PubMed](#)]
81. Masumoto S, Akimoto Y, Oike H, Kobori M. Dietary phloridzin reduces blood glucose levels and reverses Sglt1 expression in the small intestine in streptozotocin-induced diabetic mice. *J Agric Food Chem.* 2009;57:4651–6. [[PubMed](#)]
82. Matchett M.D, Compton K.A, Kondo M, Neto C.C, Hurta R. A. R. Lipopolysaccharide, cranberry flavonoids and regulation of ornithine decarboxylase (ODC) and spermidine/spermine N1-acetyltransferase (SSAT) expression in H-ras transformed cells. *FASEB J.* 2005;19(A825)
83. Matchett M.D, MacKinnon S.L, Sweeney M.I, Gottschall-Pass K.T, Hurta R. A. R. Inhibition of matrix metalloproteinase activity in DU145 human prostate cancer cells by flavonoids from lowbush blueberry (*Vaccinium angustifolium*): Possible roles for protein kinase C and mitogen activated protein kinase mediated events. *J Nutr Biochem.* 2006;17:117–25. [[PubMed](#)]
84. Matsushima M, Suzuki T, Masui A, editors. Growth inhibitory action of cranberry on *Helicobacter pylori*. *J Gastroenterol Hepatol.* 2008 2:S175–80. [[PubMed](#)]
85. McKay D.L, Blumberg J.B. Cranberries (*Vaccinium macrocarpon*) and cardiovascular disease risk factors. *Nutr Rev.* 2007;65:490–502. [[PubMed](#)]
86. McMurdo M.E, Argo I, Phillips G, Daly F, Davey P. Cranberry or trimethoprim for the prevention of recurrent urinary tract infections? A randomized controlled trial in older women. *J Antimicrob Chemother.* 2009;63:389–95. [[PMC free article](#)] [[PubMed](#)]
87. McMurdo M. E. T, Bissett L.Y, Price R. J. G, Phillips G, Crombie I.K. Does ingestion of cranberry juice reduce symptomatic urinary tract infections in older people in hospital? A double-blind, placebo-controlled trial. *Age Ageing.* 2005;34:256–61. [[PubMed](#)]
88. Morrow D. M. P, Fitzsimmons P. E. E, Chopra M, McGlynn H. Dietary supplementation with the antitumor promoter quercetin: Its effects on matrix metalloproteinase gene regulation. *Mutat Res.* 2001;480:269–76. [[PubMed](#)]
89. MotherLindas.com.<http://www.motherlindas.com/cranberries.htm>"Cranberries: America's Bouncing Berry".(accessed October 5, 2009).
90. Mullen W, Marks S.C, Crozier A. Evaluation of phenolic compounds in commercial fruit juices and fruit drinks. *J Agric Food Chem.* 2007;55:3148–57. [[PubMed](#)]
91. Murphy B.T, MacKinnon S.L, Yan X, Hammond G.B, Vaisberg A.J, Neto C.C. Identification of triterpene hydroxycinnamates with in vitro antitumor activity from whole cranberry fruit (*Vaccinium macrocarpon*). *J Agric Food Chem.* 2003;51:3541–5. [[PubMed](#)]
92. Neto C.C. Identification of triterpene hydroxycinnamates with in vitro cancer and vascular diseases. *J Agric Food Chem.* 2003;50:5844–9.
93. Neto C.C. Cranberry and blueberry: Evidence for protective effects against cancer and vascular diseases. *Mol Nutr Food Res.* 2007a;51:652–64. [[PubMed](#)]
94. Neto C.C. Cranberry and its phytochemicals: A review of in vitro anticancer studies. *J Nutr.* 2007b;137:186S–93S. [[PubMed](#)]
95. Neto C.C, Krueger C.G, Lamoureaux T.L, editors. MALDI-TOF MS characterization of proanthocyanidins from cranberry fruit (*Vaccinium macrocarpon*) that inhibit tumor cell

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

- growth and matrix metalloproteinase expression in vitro. *J Sci Food Agric.* 2006;86:18–25.
96. Novotny L, Vachalkova A, Biggs D. Ursolic acid: An anti-tumorigenic and chemopreventive activity minireview. *Neoplasma.* 2001;48:241–6. [[PubMed](#)]
 97. Ofek I, Goldhar J, Zafriri D, Lis H, Adar R, Sharon N. Anti-*Escherichia* adhesion activity of cranberry and blueberry juices. *N Eng J Med.* 1991;324:1599. [[PubMed](#)]
 98. Ohnishi R, Ito H, Kasajima N, editors. Urinary excretion of anthocyanins in humans after cranberry juice ingestion. *Biosci Biotechnol Biochem.* 2006;70:1681–7. [[PubMed](#)]
 99. O’Leary K, Pascual-Teresa S. de, Needs P.W, Bao Y.P, O’Brien N.M, Williamson G. Effect of flavonoids and Vitamin E on cyclooxygenase-2 (COX-2) transcription. *Mutat Res.* 2004;551:245–4. [[PubMed](#)]
 100. Pedersen C.B, Kyle J, Jenkinson A.M, Gardner P.T, McPhail D.B, Duthie G.G. Effects of blueberry and cranberry juice consumption on the plasma antioxidant capacity of healthy female volunteers. *Eur J Clin Nutr.* 2000;54:405–8. [[PubMed](#)]
 101. Pinzon-Arango P.A, Liu Y, Camesano T.A. Role of cranberry on bacterial adhesion forces and implications for *Escherichia coli*-uroepithelial cell attachment. *J Med Food.* 2009;12:259–70. [[PubMed](#)]
 102. Prasain J, Jones K, Moore R, editors. Effect of cranberry juice concentrate on chemically-induced bladder cancers. *Oncol Rep.* 2008;19:1565–70. [[PMC free article](#)] [[PubMed](#)]
 103. Prior R.L, Lazarus S.A, Cao G, Muccitelli H, Hammerstone J.F. Identification of procyanidins and anthocyanins in blueberries and cranberries (*Vaccinium* spp.) using high-performance liquid chromatography/mass spectrometry. *J Agric Food Chem.* 2001;49:1270–6. [[PubMed](#)]
 104. Pupa S.M, Menard S, Fortiand S, Tagliabue E. New insights into the role of extracellular matrix during tumor onset and progression. *J Cell Physiol.* 2002;192:259–67. [[PubMed](#)]
 105. Ranelletti F.O, Maggiano N, Serra F.G. Quercetin inhibits p21-Ras expression in human colon cancer cell lines and in primary colorectal tumors. *Int J Cancer.* 2000;85:438–45. [[PubMed](#)]
 106. Recio M.C, Giner R.M, Manez S, editors. Investigations on the steroidal anti-inflammatory activity of triterpenoids from *Diospyros leucomelas*. *Planta Med.* 1995;61:9–12. [[PubMed](#)]
 107. Richter M, Ebermann R, Marian B. Quercetin-induced apoptosis in colorectal tumor cells: Possible role of EGF receptor signaling. *Nutr Cancer.* 1999;34:88–99. [[PubMed](#)]
 108. Ringbom T, Segura L, Noreen Y, Bohlin L. Ursolic acid from *Plantago major*, a selective inhibitor of cyclooxygenase-2 catalyzed prostaglandin biosynthesis. *J Nat Prod.* 1998;61:1212–5. [[PubMed](#)]
 109. Roy S, Khanna S, Alessio H.M, editors. Antiangiogenic property of edible berries. *Free Radic Res.* 2002;36:1023–31. [[PubMed](#)]
 110. Ruel G, Couillard C. Evidences of the cardioprotective potential of fruits: The case of cranberries. *Mol Nutr Food Res.* 2007;51:692–701. [[PubMed](#)]
 111. Ruel G, Pomerleau S, Couture P, Lamarche B, Couillard C. Changes in plasma antioxidant capacity and oxidized low-density lipoprotein levels in men after short-term cranberry juice consumption. *Metabolism.* 2005;54:856–61. [[PubMed](#)]

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

112. Safayhi H, Sailer E.R. Anti-inflammatory actions of pentacyclic triterpenes. *Planta Med.* 1997;63:487–93. [[PubMed](#)]
113. Schmidt B.M, Erdman J.W, Lila M.A. Differential effects of blueberry proanthocyanidins on androgen sensitive and insensitive human prostate cancer cell lines. *Cancer Lett.* 2006;231:240–6. [[PubMed](#)]
114. Seeram N.P. Berries. Maryland Heights, MO: Academic Press/Elsevier; Nutritional Oncology. 2006 Chapter 37.
115. Seeram N.P, Adams L.S, Hardy M.L, Heber D. Total cranberry extract versus its phytochemical constituents: Antiproliferative and synergistic effects against human tumor cell lines. *J Agric Food Chem.* 2004;52:2512–7. [[PubMed](#)]
116. Seeram N.P, Momin R.A, Nair M.G, Bourquin L.D. Cyclooxygenase inhibitory and antioxidant cyanidin glycosides in cherries and berries. *Phytomed.* 2001;8:362–9. [[PubMed](#)]
117. Seeram N.P, Zhang Y, Nair M.G. Inhibition of proliferation of human cancer cells and cyclooxygenase enzymes by anthocyanidins and catechins. *Nutr Cancer.* 2003;46:101–6. [[PubMed](#)]
118. Sesink A. L. A, O’Leary K.A, Hollman P. C. H. Quercetin glucuronides but not glucosides are present in human plasma after consumption of quercetin-3-glucoside or quercetin-4'-glucoside. *J Nutr.* 2001;131:1938–41. [[PubMed](#)]
119. Sheng H, Shao J, Kirkland S.C, editors. Inhibition of human colon cancer cell growth by selective inhibition of cyclooxygenase-2. *J Clin Invest.* 1997;99:2254–9. [[PMC free article](#)] [[PubMed](#)]
120. Shishodia S, Majumdar S, Banerjee S, Aggarwal B.B. Ursolic acid inhibits nuclearfactor-KB activation induced by carcinogenic agents through suppression of IKBa kinase and p65 phosphorylation: Correlation with down-regulation of cyclooxygenase 2, matrix metalloproteinase 9, and cyclin D1. *Cancer Res.* 2003;63:4375–83. [[PubMed](#)]
121. Singh A.J, Singh R.K, Kim K.K, editors. Cranberry proanthocyanidins are cytotoxic to human cancer cells and sensitize platinum-resistant ovarian cancer cells to paraplatin. *Phytother Res.* 2009;23:1066–74. [[PMC free article](#)] [[PubMed](#)]
122. Singletary K, Meline B. Effect of grape seed proanthocyanidins on colon aberrant crypts and breast tumors in a rat dual-organ tumor model. *Nutr Cancer.* 2001;39:252–8. [[PubMed](#)]
123. Sobota A.E. Inhibition of bacterial adherence by cranberry juice: Potential use for the treatment of urinary tract infections. *J Urol.* 1984;131:1013–6. [[PubMed](#)]
124. Steinberg D. The LDL modification hypothesis of atherogenesis: An update. *J Lipid Res.* 2009 50:S376–81. [[PMC free article](#)] [[PubMed](#)]
125. Stothers L. A randomized trial to evaluate effectiveness and cost effectiveness of naturopathic cranberry products as prophylaxis against urinary tract infection in women. *Can J Urol.* 2002;9:1558–62. [[PubMed](#)]
126. Subbaramaiah K, Michaluart P, Sporn M.B, Dannenberg A.J. Ursolic acid inhibits cyclooxygenase-2 transcription in human mammary epithelial cells. *Cancer Res.* 2000;60:2399–404. [[PubMed](#)]
127. Sun J, Liu R.H. Cranberry phytochemical extracts induce cell cycle arrest and apoptosis in human MCF-7 breast cancer cells. *Cancer Lett.* 2006;241:124–34. [[PubMed](#)]

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

128. Turner A, Chen S.N, Joike M.K, Pendland S.L, Pauli G.F, Farnsworth N.R. Inhibition of uropathogenic *Escherichia coli* by cranberry juice: A new antiadherence assay. *J Agric Food Chem.* 2005;53:8940–7. [\[PubMed\]](#)
129. Turner A, Chen S.N, Nikolic D, van Breemen R, Farnsworth N, Pauli G. Coumaroyl iridoids and a depside from cranberry (*Vaccinium macrocarpon*). *J Nat Prod.* 2007;70:253–8. [\[PMC free article\]](#) [\[PubMed\]](#)
130. Uemura N, Okamoto S, Yamamoto S, Matsumura N, Yamaguchi S, Yamakido M. *Helicobacter pylori* infection and the development of gastric cancer. *N Engl J Med.* 2001;345:784–89. [\[PubMed\]](#)
131. USDA. Database for the Flavonoid Content of Selected Foods, Release 2.1. 2007. <http://www.ars.usda.gov/nutrientdata> (accessed December 14, 2010)
132. USDA. Database for the Proanthocyanidin Content of Selected Foods. 2004. <http://www.nal.usda.gov/fnic/foodcomp/Data/PA/PA.pdf> (accessed December 14, 2010)
133. USDA/Economic Research Service. Food Availability (Per Capita) Data System. <http://www.ers.usda.gov/Data/FoodConsumption> (accessed February 16 2010).
134. Valentova K, Stejskal D, Bednar P, editors. Biosafety, antioxidant status, and metabolites in urine after consumption of dried cranberry juice in healthy women: A pilot double-blind placebo-controlled trial. *J Agric Food Chem.* 2007;55:3217–24. [\[PubMed\]](#)
135. Vinson J.A, Bose P, Proch J, AlKharrat H, Samman N. Cranberries and cranberry products: Powerful in vitro, ex vivo, and in vivo sources of antioxidants. *J Agric Food Chem.* 2008;56:5884–91. [\[PubMed\]](#)
136. Vinson J.A, Su X, Zubik L, Bose P. Phenol antioxidant quantity and quality in foods: Fruits. *J Agric Food Chem.* 2001;49:5315–21. [\[PubMed\]](#)
137. Volate S.R, Davenport D.M, Muga S.J, Wargovich M.J. Modulation of aberrant crypt foci and apoptosis by dietary herbal supplements (quercetin, curcumin, silymarin, ginseng, and rutin). *Carcinogenesis.* 2005;26:1450–6. [\[PubMed\]](#)
138. Vorsa N, Howell A.B. Structure and genetic variation of cranberry proanthocyanidins that inhibit adherence of uropathogenic P-fimbriated *E. coli*. Washington, DC: ACS Books.; In *Food Factors in Health Promotion and Disease Prevention, Symposium Series, 851.* 2003
139. Vvedenskaya I.O, Rosen R.T, Guido J.E, Russell D.J, Mills K.A, Vorsa N. Characterization of flavonols in cranberry (*Vaccinium macrocarpon*) powder. *J Agric Food Chem.* 2004;52:188–95. [\[PubMed\]](#)
140. Vvedenskaya I.O, Vorsa N. Flavonoid composition over fruit development and maturation in American cranberry, *Vaccinium macrocarpon* Ait. *Plant Sci.* 2004;167:1043–54.
141. Walker E.B, Barney D.P, Mickelson J.N, Walton R.J, Mickelson, Jr. R.A. Cranberry concentrate:UTI prophylaxis. *J Fam Pract.* 1997;45:167–8. [\[PubMed\]](#)
142. Wang M, Li J, Shao Y, editors. Champaign, IL: AOCS Press.; *Antioxidative and Cytotoxic Components of Highbush Blueberry (Vaccinium corymbosum L.) Phytochem Phytopharmaceus.* 2000
143. Wang S.Y, Stretch A.W. Antioxidant capacity in cranberry is influenced by cultivar and storage temperature. *J Agric Food Chem.* 2001;49:969–74. [\[PubMed\]](#)
144. Wang Y, Catana F, Yang Y, Roderick R, Breemen R. B. van. An LC-MS method for analyzing total resveratrol in grape juice, cranberry juice, and in wine. *J Agric Food Chem.* 2002;50:431–5. [\[PubMed\]](#)

Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition.

145. Wilson T, Meyers S.L, Singh A.P, Limburg P.J, Vorsa N. Favorable glycemic response of type 2 diabetics to low-calorie cranberry juice. *J Food Sci.* 2008;73:H241–5. [[PubMed](#)]
146. Wing D.A, Rumney P.J, Preslicka C.W, Chung J.H. Daily cranberry juice for the prevention of asymptomatic bacteriuria in pregnancy: A randomized, controlled pilot study. *J Urol.* 2008;180:1367–72. [[PMC free article](#)] [[PubMed](#)]
147. Wu X, Beecher G.R, Holden J.M, Haytowitz D.B, Gebhardt S.E, Prior R.I. Concentrations of anthocyanins in common foods in the United States and estimation of normal consumption. *J Agric Food Chem.* 2006;54:4069–75. [[PubMed](#)]
148. Wu X, Prior R.J. Systematic identification and characterization of anthocyanins by HPLC-ESI-MS/ MS in common foods in the United States: Fruits and berries. *J Agric Food Chem.* 2005;53:2589–99. [[PubMed](#)]
149. Yan X, Murphy B, Hammond G.B, Vinson J.A, Neto C.C. Antioxidant activities and antitumor screening of extracts from cranberry fruit (*Vaccinium macrocarpon*). *J Agric Food Chem.* 2002;20:5844–9. [[PubMed](#)]
150. Yang G.-Y, Liao J, Kim K, Yurkow E.J, Yang C.S. Inhibition of growth and induction of apoptosis in human cancer cell lines by tea polyphenols. *Carcinogenesis.* 1998;19:611–6. [[PubMed](#)]
151. Zafriri D, Ofek I, Pocino A.R, Sharon N. Inhibitory activity of cranberry juice on adherence of type 1 and type P-fimbriated *Escherichia coli* to eucaryotic cells. *Antimicrob Agents Chemother.* 1989;33:92–8. [[PMC free article](#)] [[PubMed](#)]
152. Zhang K, Zuo Y. GC-MS Determination of flavonoids and phenolic and benzoic acids in human plasma after consumption of cranberry juice. *J Agric Food Chem.* 2004;52:222–7. [[PubMed](#)]
153. Zhang L, Ma J, Pan K, Go V.L, Chen J, You W.C. Efficacy of cranberry juice on *Helicobacter pylori* infection: A double-blind randomized placebo-controlled trial. *Helicobacter.* 2005;10:139–45. [[PubMed](#)]
154. Zhao Y, editor. Boca Raton, FL: CRC Press; *Berry Fruit: Value-Added Products for Health Promotion.* 2007
155. Zheng W, Wang S.Y. Oxygen radical absorbing capacity of phenolics in blueberries, cranberries, chokeberries, and lingonberries. *J Agric Food Chem.* 2003;51:502–9. [[PubMed](#)]
156. Zuo Y, Wang C, Zhan J. Separation, characterization, and quantitation of benzoic and phenolic antioxidants in American cranberry fruit by GC-MS. *J Agric Food Chem.* 2002;50:3789–94. [[PubMed](#)]